



Ref. 302004

Last update 15/04/2022

### Description & Benefits

Complete feed in pellet form for the maintenance of ornamental poultry outside the breeding season:

**Pheasant, partridge, turkey, bustard.**

Good fiber, vitamin and mineral content.



### Feeding Instructions

Supply with fresh clean water.

**Warning:** This feed must not be distributed to animals intended for human consumption.

### Composition

Wheat, alfalfa, corn, soybean meal, oats, wheat bran, peas, dicalcium phosphate, rapeseed oil, vitamins and trace elements mix, salt, yeast, fenugreek, methionine, antifungal preservatives, polyphenol and tocopherols extracts.

### Packaging & Preservation

Bag of 25 kg  
Extrudates 4 x 8 mm  
Store away from light, in a cool and dry place.



## GALLIFORMS DIET MAINTENANCE



### Analysis

#### KEY NUTRIENTS

Crude protein	16.60 %
Crude fat	5 %
Crude fibre	7.70 %
Crude ash	7.20 %
Moisture	9 %
NDF	20.80 %
ADF	9.80 %
ADL	2.20 %
Starch	35.10 %
NFE	54.10 %

#### MINERALS

Calcium	1.20 %
Phosphorus	0.80 %
Sodium	1.70 %
Potassium	10 %
Magnesium	1.60 %

#### TRACE ELEMENTS

Iron	200 mg/kg
Copper	20 mg/kg
Manganese	60 mg/kg
Zinc	100 mg/kg
Cobalt	0.20 mg/kg
Iodine	0.50 mg/kg
Selenium	0.50 mg/kg

#### AMINO ACIDS

Arginine	9.80 g/kg
Lysine	7.50 g/kg
Methionine	4.90 g/kg
Cystine	3 g/kg
Tryptophan	1.90 g/kg
Threonine	5.70 g/kg

#### VITAMINS<sup>1</sup>

Vitamin A	16500 UI
Vitamin D3	3150 UI
Vitamin E	300 mg/kg
Menadione (K3)	50 mg/kg
Thiamin (B1)	28 mg/kg
Riboflavin (B2)	12 mg/kg
Nicotinic acid (B3)	50 mg/kg
Pantothenic adic (B5)	25 mg/kg
Pyridoxin (B6)	10 mg/kg
Biotin (B8)	0.30 mg/kg
Folic acid (B9)	6.50 mg/kg
Cobalamin (B12)	50 mg/kg
Choline	2000 mg/kg

#### ENERGY

Gross energy	16.70 MJ/kg
Metabolisable energy <sup>2</sup>	10.20 MJ/kg

Hereabove values correspond to calculated average values. These values may vary due to the natural variability of raw materials and manufacturing.

1 : Values for vitamins correspond to added quantities during manufacturing.

2 : Metabolisable energy for ruminants according to the INRA 2018 system.